

**DRUG FACTS**

Active ingredient(s)

Purpose

**Use(s)**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have**  
**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are**  
**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**  
**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**

**Keep out of reach of children**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- store between 20° to 25°C (68° to 77°F)
- **do not use if imprinted foil inner seal on bottle is broken or missing**

**Drug Facts**

**Inactive ingredients**  
colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide